

Accepted Manuscript

Letter to the Editor

Studies of macrophage therapy for cirrhosis – from mice to men

James A Thomas, Prakash Ramachandran, Stuart J Forbes

PII: S0168-8278(18)30002-3

DOI: <https://doi.org/10.1016/j.jhep.2017.11.043>

Reference: JHEPAT 6809


To appear in: *Journal of Hepatology*

Received Date: 7 November 2017

Accepted Date: 28 November 2017

Please cite this article as: Thomas, J.A., Ramachandran, P., Forbes, S.J., Studies of macrophage therapy for cirrhosis – from mice to men, *Journal of Hepatology* (2018), doi: <https://doi.org/10.1016/j.jhep.2017.11.043>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Title page:

Letter to Editor re Ma et al (Cytotherapy with M1-polarized macrophages ameliorates liver fibrosis by modulating immune microenvironment in mice. J Hepatol. 2017 Oct;67(4):770-779).

Authors:

Dr James A Thomas – 1) The Prince Charles Hospital. Brisbane. Queensland,
2) University of Queensland. Australia

Dr Prakash Ramachandran – University of Edinburgh. UK

Prof Stuart J Forbes – University of Edinburgh. UK

Title: Studies of macrophage therapy for cirrhosis – from mice to men

Studies of macrophage therapy for cirrhosis – from mice to men

We read with interest the recently published paper by Ma et al. (1) in which they examined the effects of macrophage infusion in experimental liver fibrosis. The beneficial effects on measures of liver scarring and regeneration following the delivery of bone marrow derived macrophages (BMDMs) suggest clinical potential.

BMDMs injected during iterative liver injury resulted in the hepatic engraftment of relatively few donor cells. Consistent with our previously reported findings (2), macrophage administration resulted in modulation of the host immune response with upregulation of antifibrotic matrix metalloproteinases, apoptosis of scar producing myofibroblasts and subsequent reduction in liver fibrosis. The injection of unpolarized (naïve) BMDMs mediated these effects as did those previously stimulated *in vitro* to a classical "M1" phenotype. In contrast, BMDMs activated to the alternative "M2" phenotype were not beneficial.

Our group's work characterizing the "restorative macrophage" during the spontaneous regression of fibrosis after liver injury, has shown that these phagocytic Ly6C^{lo} cells do not conform to the traditional M1-M2 paradigm (3). The hepatic macrophage population extracted from recipient liver after BMDM therapy in this paper appeared to have a comparable phenotype. This could indicate that BMDM therapy is able to harness this endogenous repair process, critically, in the context of ongoing liver injury. The demonstrated upregulation of macrophage chemokines (CCL-2 and -3) is consistent with the hepatic recruitment of host cells. Amplification of the effects of the initial cell injection through paracrine signaling is an attractive concept to explain the enduring changes found weeks after BMDM delivery. However, this does not exclude a significant contribution to the increased numbers of hepatic F4/80+ macrophages from proliferation of resident hepatic macrophages in addition to recruitment (4).

Ma et al. (1) also demonstrated that BMDM infusion increased the number of activated hepatic NK cells, with the potential to promote myofibroblast apoptosis. Whether this NK cell enrichment reflects a direct effect of the infused BMDMs or a consequence of the modulation in host macrophage phenotype remains to be elucidated. Further studies, administering BMDMs deficient in specific chemokines could provide further mechanistic insights.

We have reported increased serum albumin levels following BMDM therapy (2). This finding has also been recapitulated by Ma et al. (1). The increase in Ki67+ hepatocytes identifies hepatocyte proliferation as a regenerative mechanism after BMDM delivery. Our work exploring BMDM therapy mediated regeneration in both the injured (2) and uninjured (5) liver has also indicated a role for the liver progenitor cell (LPC) mitogen TWEAK. This cytokine specifically activates the LPC compartment thereby increasing liver cell mass. Evaluation of this axis following injection of differentially polarized macrophages would be informative from biological and translational perspectives.

The authors also tested BMDM therapy in cholestatic fibrosis due to bile duct ligation. The antifibrotic effect was found to be of a similar magnitude to that detected in the carbon tetrachloride (CCl₄) induced model of hepatic injury. The mechanistic studies reported in this article were carried out exclusively in the CCl₄ model. We have previously demonstrated fundamental differences in the spatial and in turn functional characteristics of biliary versus hepatic fibrosis and regeneration (6). Therefore, further exploration of the processes underlying the effects observed in this cholestatic model will be of great interest.

Following our initial report (2), we have sought to test the translational potential of BMDM therapy. Human macrophages injected into mice with experimental liver fibrosis showed similar positive effects on liver fibrosis and function (7). Building on this infrastructure, our group has validated and scaled-up a Good Manufacturing Practice-compliant process for deriving human macrophages for therapy (8). A first in man study of autologous macrophage infusion in cirrhosis (ISRCTN 10368050) has already commenced to determine whether these promising pre-clinical findings can yield effective treatment for patients with cirrhosis.

References:

1. Ma PF, Gao CC, Yi J, Zhao JL, Liang SQ, Zhao Y, Ye YC, Bai J, Zheng QJ, Dou KF, Han H, Qin HY. Cytotherapy with M1-polarized macrophages ameliorates liver fibrosis by modulating immune microenvironment in mice. *J Hepatol*. 2017 Oct;67(4):770-779.
2. Thomas JA, Pope C, Wojtacha D, Robson AJ, Gordon-Walker TT, Hartland S, Ramachandran P, Van Deemter M, Hume DA, Iredale JP, Forbes SJ. Macrophage therapy for murine liver fibrosis recruits host effector cells improving fibrosis, regeneration, and function. *Hepatology*. 2011 Jun;53(6):2003-15.
3. Ramachandran P, Pellicoro A, Vernon MA, Boulter L, Aucott RL, Ali A, Hartland SN, Snowden VK, Cappon A, Gordon-Walker TT, Williams MJ, Dunbar DR, Manning JR, van Rooijen N, Fallowfield JA, Forbes SJ, Iredale JP. Differential Ly-6C expression identifies the recruited macrophage phenotype, which orchestrates the regression of murine liver fibrosis. *Proc Natl Acad Sci U S A*. 2012 Nov 13;109(46):3186-95.
4. Stutchfield BM, Antoine DJ, Mackinnon AC, Gow DJ, Bain CC, Hawley CA, Hughes MJ, Francis B, Wojtacha D, Man TY, Dear JW, Devey LR, Mowat AM, Pollard JW, Park BK, Jenkins SJ, Simpson KJ, Hume DA, Wigmore SJ, Forbes SJ. CSF1 Restores Innate Immunity After Liver Injury in Mice and Serum Levels Indicate Outcomes of Patients With Acute Liver Failure. *Gastroenterology*. 2015 Dec;149(7):1896-1909.
5. Bird TG, Lu WY, Boulter L, Gordon-Keylock S, Ridgway RA, Williams MJ, Taube J, Thomas JA, Wojtacha D, Gambardella A, Sansom OJ, Iredale JP, Forbes SJ. Bone marrow injection stimulates hepatic ductular reactions in the absence of injury via macrophage-mediated TWEAK signaling. *Proc Natl Acad Sci U S A*. 2013 Apr 16;110(16):6542-7.
6. Boulter L, Govaere O, Bird TG, Radulescu S, Ramachandran P, Pellicoro A, Ridgway RA, Seo SS, Spee B, Van Rooijen N, Sansom OJ, Iredale JP, Lowell S, Roskams T, Forbes SJ. Macrophage-derived Wnt opposes Notch signaling to specify hepatic progenitor cell fate in chronic liver disease. *Nat Med*. 2012 Mar 4;18(4):572-9.
7. Moore JK, Mackinnon AC, Wojtacha D, Pope C, Fraser AR, Burgoyne P, Bailey L, Pass C, Atkinson A, McGowan NW, Manson L, Turner ML, Campbell JD, Forbes SJ. Phenotypic and functional characterization of macrophages with therapeutic potential generated from human cirrhotic monocytes in a cohort study. *Cytotherapy*. 2015 Nov;17(11):1604-16.
8. Fraser AR, Pass C, Burgoyne P, Atkinson A, Bailey L, Laurie A, W A McGowan N, Hamid A, Moore JK, Dwyer BJ, Turner ML, Forbes SJ, Campbell JDM. Development, functional characterization and validation of methodology for GMP-compliant manufacture of phagocytic macrophages: A novel cellular therapeutic for liver cirrhosis. *Cytotherapy*. 2017 Sep;19(9):1113-1124.